

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of
GOERLITZER, J., et. al.

Examiner: Paviglianiti, Anthony Joseph

Application No.: **10/789,019**

Art Unit: 1626

Filed: **February 27, 2004**

Title: **Diarylcycloalkyl derivatives, process for
their preparation and their use as
pharmaceuticals**

TELEFAX CERTIFICATE

I hereby certify that this correspondence is
being transmitted via facsimile to the
Commissioner for Patents, Alexandria,
VA 22313, on

August 22, 2005
Date of Transmission

Brian Pritchett
Signature

Brian Pritchett

Terminal Disclaimer

Commissioner for Patents
P. O. Box 1450
Alexandria, VA 22313-1450

**RECEIVED
CENTRAL FAX CENTER**

AUG 22 2005

Your petitioner, Aventis Pharma Deutschland GmbH, a corporation having offices in Industriepark Hoechst, Gebaude K801, Frankfurt Am Main, Germany, represents that it is the assignee of Application Serial No. 10/789,019, filed on February 27, 2004, by virtue of an assignment to Aventis Pharma Deutschland GmbH which was recorded on June 11, 2004, set out in reel 014722 and frame 0060.


Your petitioner, Aventis Pharma Deutschland GmbH, hereby disclaims the terminal part of any patent granted on the above-identified application, which would extend beyond the expiration date of the patent corresponding to USSN 10/789,281, which is assigned to and commonly owned by Aventis Pharma Deutschland GmbH and hereby agrees that any patent so granted on the above-identified application shall be enforceable only for and during such period that the legal title to said patent shall be the same as the legal title to United States Patent Application Serial No. 10/789,281, this agreement to run with any patent granted on the above-identified application and to be binding upon the grantee, its successors or assigns. Such disclaimer does not, however, mean that petitioner hereby waives or disclaims any right to an extension of the term of any patent issuing from application serial number 10/789,019, under 35 U.S.C. § 156, pursuant to (a) Title II of Pub. L. No. 98-417, enacted on September 24, 1984, and entitled the "Drug Price Competition and Patent Term Restoration Act of 1984", (b) Pub. 98-417, enacted on September 24, 1984, and entitled the "Drug Price Competition and Patent Term Restoration Act of 1984", (b) Pub. L. No. 100-670, enacted November 16, 1988 and entitled "Patent Term Extension for Animal

Drug Products", or (c) any other subsequently enacted Public Law or Act which provides for the extension of the term of a patent.

Such disclaimer does not disclaim any terminal part of any patent granted on the above-identified application prior to the expiration date of the full statutory term, in the event that it later expires for failure to pay a maintenance fee, is held unenforceable, is found invalid, is statutorily disclaimed in whole or terminally disclaimed under 37 CFR 1.321(a), has all claims cancelled by a reexamination certificate, or is otherwise terminated prior to the expiration of its statutory term as presently shortened by any terminal disclaimer, except for the separation of legal title as stated above.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon. Please charge Deposit Account No. 18-1982 in the amount of \$110.00. Two duplicate copies of this sheet are enclosed. The Commissioner is authorized to charge any fees under 37 CFR 1.16 - 1.21 or credit any overpayment to Deposit Account No. 18-1982.

Respectfully submitted,



Barbara E. Kurys, Reg. No. 34,650


Attorney/Agent For Applicant

Aventis Pharmaceuticals Inc.
Patent Department
Route 202/205
P.O. Box 6800
Bridgewater, NJ 08807-0800
Telephone (908) 231-2965
Telefax (908) 231-2626

DEAV2003/0017 US NP

Facsimile Transmission

This company uses RightFAX® fax server software from RightFAX, Inc.



From: Name: USPTO ASSIGNMENT DIVISION
Fax Number:
Voice Phone: 703-308-9723

To: Name: AVENTIS PHARMACEUTICALS INC.
Company: BARBARA E. KURYS
Fax Number: 19082312626
Voice Phone:

Fax Notes:

Pg#	Description
1	Cover Page
2	375.TXT
4	Document 1, Batch 319949

PTAS FAX PROCESSING

for ACTION DUE file D9AU2003/0017-05-NP

DUE DATE	
EXTD BY	Jes 6/15/04
ATTY.	B&K/R.Wingefeld

Date and time of transmission: Monday, June 14, 2004 2:36:08 PM
Number of pages including this cover sheet: 04

A RightFAX® Communicated Document



UNITED STATES DEPARTMENT OF COMMERCE
Patent and Trademark Office
ASSISTANT SECRETARY AND COMMISSIONER
OF PATENTS AND TRADEMARKS
Washington, D.C. 20231



700090914A

JUNE 14, 2004

PTAS

AVENTIS PHARMACEUTICALS INC.
BARBARA E. KURYS
ROUTE 202-206
P.O. BOX 6800
BRIDGEWATER, NJ 08807-00800

UNITED STATES PATENT AND TRADEMARK OFFICE
NOTICE OF RECORDATION OF ASSIGNMENT DOCUMENT

THE ENCLOSED DOCUMENT HAS BEEN RECORDED BY THE ASSIGNMENT DIVISION OF THE U.S. PATENT AND TRADEMARK OFFICE. A COMPLETE MICROFILM COPY IS AVAILABLE AT THE ASSIGNMENT SEARCH ROOM ON THE REEL AND FRAME NUMBER REFERENCED BELOW.

PLEASE REVIEW ALL INFORMATION CONTAINED ON THIS NOTICE. THE INFORMATION CONTAINED ON THIS RECORDATION NOTICE REFLECTS THE DATA PRESENT IN THE PATENT AND TRADEMARK ASSIGNMENT SYSTEM. IF YOU SHOULD FIND ANY ERRORS OR HAVE QUESTIONS CONCERNING THIS NOTICE, YOU MAY CONTACT THE EMPLOYEE WHOSE NAME APPEARS ON THIS NOTICE AT 703-308-9723. PLEASE SEND REQUEST FOR CORRECTION TO: U.S. PATENT AND TRADEMARK OFFICE, ASSIGNMENT DIVISION, BOX ASSIGNMENTS, CG-4, 1213 JEFFERSON DAVIS HWY, SUITE 320, WASHINGTON, D.C. 20231.

RECORDATION DATE: 06/11/2004

REEL/FRAME: 014722/0060

NUMBER OF PAGES: 2

BRIEF: ASSIGNMENT OF ASSIGNOR'S INTEREST (SEE DOCUMENT FOR DETAILS).

ASSIGNOR:

GOERLITZER, JOCHEN

DOC DATE: 05/04/2004

ASSIGNOR:

GLOMBIK, HEINER

DOC DATE: 05/06/2004

ASSIGNOR:

FALK, EUGEN

DOC DATE: 05/19/2004

ASSIGNOR:

GRETZKE, DIRK

DOC DATE: 05/11/2004

ASSIGNOR:

KEIL, STEFANIE

DOC DATE: 05/04/2004

ASSIGNOR:

SCHAEFER, HANS-LUDWIG

DOC DATE: 05/04/2004

014722/0060 PAGE 2

ASSIGNOR:
STAPPER, CHRISTIAN

DOC DATE: 05/14/2004

ASSIGNOR:
WENDLER, WOLFGANG

DOC DATE: 05/14/2004

ASSIGNEE:
AVENTIS PHARMA DEUTSCHLAND GMBH
FRANKFURT AM MAIN, FED REP
GERMANY
D-69529

SERIAL NUMBER: 10789019

FILING DATE: 02/27/2004

PATENT NUMBER:

ISSUE DATE:

TITLE: DIARYLCYCLOALKYL DERIVATIVES, PROCESS FOR THEIR PREPARATION AND
THEIR USE AS PHARMACEUTICALS

SHAREILL COLES, EXAMINER
ASSIGNMENT DIVISION
OFFICE OF PUBLIC RECORDS

UPT AUG. 22. 2005 3:43PM
JUN 11 2004 7:20AM

CAVENTIS US PAT DEPT 4/4

High FAX

NO. 7674
NO. 4034 r. 1

RECEIVED
CENTRAL FAX CENTER

AUG 22 2005

TQ: US USPTO AS 10661172004
780690914

FORM PTO-1695
1-31-92
(modified)

RECORDATION FORM COVER SHEET
PATENTS ONLY

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE

I hereby certify that this correspondence is being transmitted via facsimile to the Commissioner for Patents, Alexandria, VA 22313-1450, Patent and Trademark Assignment System, at (703) 306-6983, on
Date June 11, 2004

Signature Barbara E. Kurys

To the Commissioner of Patents. Please record the attached original documents or copy thereof.

1. Name of conveying party(ies):

- 1) Jochen Goerlitzer
- 2) Heiner Glombik
- 3) Eugen Falk
- 4) Dirk Gretzke

- 5) Stefanie Keil
- 6) Hans-Ludwig Schaefer
- 7) Christian Stapper
- 8) Wolfgang Wendler

Additional name(s) of conveying party(ies) attached? ☐ YES ☒ NO

2. Name and address of receiving party(ies):

Name: **Aventis Pharm Deutschland GmbH**
Internal Address:
Street Address:
City: **Frankfurt am Main** City: **Germany** ZIP: **D-69529**

Additional name(s) and address(es) attached? ☐ YES ☒ NO

3. Nature of Conveyance:

- ☒ Assignment ☐ Merger
- ☐ Security Agreement ☐ Change of Name
- ☐ Other

Execution Date: 1) May 4, 2004; 2) May 6, 2004; 3) May 19, 2004; 4) May 11, 2004; 5) May 4, 2004; 6) May 4, 2004; 7) May 14, 2004; 8) May 14, 2004

4. Application number(s) or patent number(s):

If this document is being filed together with a new application, the execution date of the application is:

A. Patent Application No.(s): **10/789,019**

B. Patent No.(s):

Additional numbers attached? ☐ YES ☒ NO

5. Name and address of party to whom correspondence concerning document should be mailed:

Name: **Barbara E. Kurys**
Internal Address: **Aventis Pharmaceuticals Inc.**
Street Address: **Route 202-206 / P.O. Box 6800**
City: **Bridgewater** State: **NJ** ZIP: **08807-0800**
**** FAX NUMBER: (908) 231-2626 ****
Our Reference No.: **DEAY2003/0017USNP**

6. Total number of applications and patents involved: **1**

7. Total (87 CFR 3.41): **\$40.00**

- ☐ Employed
- ☒ Authorized to be charged to deposit account

8. Deposit account number: **18-1982**

DO NOT USE THIS SPACE

9. Statement and signature.

To the best of my knowledge and belief, the foregoing information is true and correct and any attached copy is a true copy of the original document.

Barbara E. Kurys, Reg. No. 34,650
Name of Person Signing

Barbara E. Kurys
Signature

Date: 6/10/04

Total number of pages comprising cover sheet: **2**

OMB No. 0651-0011 (exp. 4/84)

Do not detach this portion
Mail documents to be recorded with required cover sheet information to:
Mail Stop Assignments
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Public burden reporting for this sample cover sheet is estimated to average about 30 minutes per document to be recorded, including time for reviewing the document and gathering the data needed, and completing and reviewing the sample cover sheet. Send comments regarding this burden estimate to the U.S. Patent and Trademark Office, Office Information Systems, PK2-10000, Alexandria, VA 22313, and to the Office of Management and Budget, Paperwork Reduction Project (0561-0011), Alexandria, VA 22313.

DEAV2003/0017 US NP

ASSIGNMENT

For valuable consideration, the receipt and adequacy of which is hereby acknowledged,

I/we

- 1) Dr. Jochen Goerlitz, Stegstrasse 60, 60594 Frankfurt am Main, Germany
- 2) Dr. Heiner Glombik, Am Lotzenwald 42, 65719 Hofheim, Germany
- 3) Dr. Eugen Falk, Völklingenweg 15, 60529 Frankfurt, Germany
- 4) Dr. Dirk Gretzke, Kaulbachstraße 57, 60596 Frankfurt, Germany
- 5) Dr. Stefanie Keil, Am Kreishaus 12, 65719 Hofheim, Germany
- 6) Dr. Hans-Ludwig Schaefer, Steingasse 7, 65239 Hochheim, Germany
- 7) Dr. Christian Stapper, Wallaustraße 53, 55118 Mainz, Germany
- 8) Dr. Wolfgang Wendler, Haintchener Str. 12a, 65618 Selters, Germany

The inventors are citizens of: 1) - 8) GERMANY .

hereby sell, assign, and transfer unto **AVENTIS PHARMA DEUTSCHLAND GMBH**, a corporation organized under the laws of **GERMANY**, located at **D-65929 Frankfurt am Main, GERMANY**,

the entire right, title and interest in and to my/our application for Letters Patent of the United States, executed concurrently herewith

Ser.-No.: 10/789,019

filed: February 27, 2004

entitled: **Diarylcyaloalkyl derivatives, process for their preparation and their use as pharmaceuticals**

and my/our entire right, title, and interest in and to all my/our inventions disclosed in said application for Letters Patent, and in and to all divisional or continuation applications that may be filed for United States Letters Patent for any of said inventions, and in and to all patents that may be granted on the foregoing applications, and I/we hereby agree, whenever requested, to communicate to said assignee, its successors and assigns, any facts known to me/us respecting said inventions, to testify in any legal proceeding, and to execute all applications or papers necessary to obtain and maintain proper patent protection on said inventions in the United States.

Inventor(s):

1 st .	<u>Jochen Goerlitz</u>	Date:	<u>5.5.2004</u>
2 nd .	<u>Heiner Glombik</u>	Date:	<u>6.5.04</u>
3 rd .	<u>Eugen Falk</u>	Date:	<u>19.5.2004</u>
4 th .	<u>Dirk Gretzke</u>	Date:	<u>11.05.2004</u>
5 th .	<u>Stefanie Keil</u>	Date:	<u>04/05/2004</u>
6 th .	<u>Hans-Ludwig Schaefer</u>	Date:	<u>14/05/2004</u>
7 th .	<u>Christian Stapper</u>	Date:	<u>4.5.2004</u>
8 th .	<u>Wolfgang Wendler</u>	Date:	<u>14.5.2004</u>

Aventis Pharma Deutschland GmbH DEAV2003/0017

Dr.WI

Diarylcycloalkyl derivatives, process for their preparation and their use as pharmaceuticals

5

The invention relates to diarylcycloalkyl derivatives and to their physiologically acceptable salts and physiologically functional derivatives.

10

Compounds of a similar structure have already been described in the prior art for the treatment of hyperlipidemia and diabetes (WO 2000/64876 and WO 03/020269).

15


The invention was based on the object of providing compounds which permit therapeutically utilizable modulation of lipid and/or carbohydrate metabolism and are thus suitable for the prevention and/or treatment of diseases such as type 2 diabetes and atherosclerosis and the diverse sequelae thereof.

20

A series of compounds which modulate the activity of PPA receptors has surprisingly been found. The compounds are suitable in particular for activating PPARalpha and PPARGgamma, it being possible for the extent of the relative activation to vary depending on the compounds.

Accordingly, the invention relates to compounds of the formula I

25



in which

Ring A is (C₃-C₈)-cycloalkanediyl or (C₃-C₈)-cycloalkenediyl, where in the

cycloalkanediyl or cycloalkenediyl rings one or more carbon atoms may be replaced by oxygen atoms;

Ring B is a) phenyl; or

5

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

10 R1 is

a) in the case ring B = a):

SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

b) in the case ring B = b):

15

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂, O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

c) in the case ring B = a) and R₄ = phenyl:

(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

20

R2 is

H or CF₃;

R4 is

a) in the case ring B = a):

phenyl;

25

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

c) in the case ring B = a) and R₁ = a):

30

(C₁-C₆)-alkyl;

R5 is

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

R3 is H or (C₁-C₆)-alkyl;

X is (C₁-C₆)-alkanediyl, where in the alkanediyl group one or more carbon atoms may be replaced by oxygen atoms;

5

Y is (C₁-C₆)-alkanediyl, where in the alkanediyl group one or more carbon atoms may be replaced by oxygen atoms;

and their physiologically acceptable salts.

10

Preference is given to compounds of the formula I in which

15

Ring A is (C₃-C₈)-cycloalkanediyl or (C₃-C₈)-cycloalkenediyl, where in the cycloalkanediyl or cycloalkenediyl rings one or more carbon atoms may be replaced by oxygen atoms;

Ring B is a) phenyl; or

20

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

R1 is a) in the case ring B = a):

25

SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂, O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

30

c) in the case ring B = a) and R4 = phenyl:
(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

R is H or CF₃;

R4 is a) in the case ring B = a):
phenyl;

5

b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

10

c) in the case ring B = a) and R1 = a):
(C₁-C₆)-alkyl;

R5 is H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

R3 is H or (C₁-C₆)-alkyl;

15

X is CH₂-O;

Y is (C₁-C₆)-alkanediyl, where in the alkanediyl group one or more carbon atoms may be replaced by oxygen atoms.

20

Preference is furthermore given to compounds of the formula I in which

Ring A is (C₃-C₈)-cycloalkanediyl in which one carbon atom may be replaced by an oxygen atom;

25

Ring B is a) phenyl; or

30

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

R1 is a) in the case ring B = a):
SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂,
O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-
alkyl;

c) in the case ring B = a) and R₄ = phenyl:

(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

10 R₂ is H or CF₃;

R₄ is a) in the case ring B = a):
phenyl;

15 b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

c) in the case ring B = a) and R₁ = a):

(C₁-C₆)-alkyl;

20

R₅ is H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;


R₃ is H or (C₁-C₆)-alkyl;

25 X is CH₂-O;

Y is CH₂-O.

Particular preference is given to compounds of the formula Ia

30



In which ring A, ring B, R1, R2, R3, R4, R5, X and Y are as defined above.

5 Particular preference is furthermore given to compounds of the formula 1a in which

R3 is H and

R5 is methyl

10

or

to compounds of the formula 1a in which

15 Ring A is (C₅-C₇)-cycloalkanediyl;

Ring B is a) phenyl; or

20

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

R1 is a) in the case ring B = a):
SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

25

b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂,

O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

5 c) in the case ring B = a) and R₄ = phenyl:
(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

R₂ is H or CF₃;

10 R₄ is a) in the case ring B = a):
phenyl;

b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

15 c) in the case ring B = a) and R₁/R₂ = a):
(C₁-C₆)-alkyl;

R₅ is methyl;

20 R₃ is H;

X is CH₂-O;

Y is CH₂-O.

25

Very particular preference is given to compounds of the formulae I and Ia

in which the central cycloalkanediyl ring is attached 1,3-cis.

30 This invention also encompasses all combinations of preferred aspects of the invention described herein.

The alkyl radicals in the substituents R₁, R₂, R₃, R₄ and R₅ may be either

straight-chain or branched.

Aryl means an aromatic carbocyclic mono- or bicyclic ring system which comprises 6 to 10 atoms in the ring or rings.

5

Heteroaryl is a mono- or bicyclic aromatic ring system having 4 to 11 ring members, in which at least one atom in the ring system is a heteroatom from the series N, O and S.

- 10 The compounds of the formula I comprise at least two centers of asymmetry and may comprise more in addition. The compounds of the formula I may therefore exist in the form of their racemates, racemic mixtures, pure enantiomers, diastereomers and mixtures of diastereomers. The present invention encompasses all these isomeric forms of the compounds of the formula I. These
- 15 isomeric forms can be obtained by known methods even if not specifically described in some cases.

- Pharmaceutically acceptable salts are, because their solubility in water is greater than that of the initial or basic compounds, particularly suitable for medical
- 20 applications. These salts must have a pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the invention are salts of inorganic acids such as hydrochloric acid, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acid, and of organic acids such as, for example, acetic acid, benzenesulfonic, benzoic, citric,
- 25 ethanesulfonic, fumaric, gluconic, glycolic, isethionic, lactic, lactobionic, maleic, malic, methanesulfonic, succinic, p-toluenesulfonic and tartaric acid. Suitable pharmaceutically acceptable basic salts are ammonium salts, alkali metal salts (such as sodium and potassium salts), alkaline earth metal salts (such as magnesium and calcium salts), and salts of trometamol (2-amino-2-
- 30 hydroxymethyl-1,3-propanediol), diethanolamine, lysine or ethylenediamine.

Salts with a pharmaceutically unacceptable anion such as, for example, trifluoroacetate likewise belong within the framework of the invention as useful

intermediates for the preparation or purification of pharmaceutically acceptable salts and/or for use in nontherapeutic, for example in vitro, applications.

5 The term "physiologically functional derivative" used herein refers to any physiologically tolerated derivative of a compound of the formula I of the invention, for example an ester, which on administration to a mammal such as, for example, a human is able to form (directly or indirectly) a compound of the formula I or an active metabolite thereof.

10 Physiologically functional derivatives also include prodrugs of the compounds of the invention, as described, for example, in H. Okada et al., Chem. Pharm. Bull. 1994, 42, 57-61. Such prodrugs can be metabolized in vivo to a compound of the invention. These prodrugs may themselves be active or not.

15 The compounds of the invention may also exist in various polymorphous forms, for example as amorphous and crystalline polymorphous forms. All polymorphous forms of the compounds of the invention belong within the framework of the invention and are a further aspect of the invention.

20 All references to "compound(s) of formula I" hereinafter refer to compound(s) of the formula I as described above, and their salts, solvates and physiologically functional derivatives as described herein.

Use

25

This invention relates further to the use of compounds of the formula I and their pharmaceutical compositions as PPAR ligands. The PPAR ligands of the invention are suitable as modulators of PPAR activity.

30 Peroxisome proliferator-activated receptors (PPAR) are transcription factors which can be activated by ligands and belong to the class of nuclear hormone receptors. There are three PPAR isoforms, PPARalpha, PPARgamma and PPARdelta, which are encoded by different genes (Peroxisome proliferator-activated receptor

(PPAR): structure, mechanisms of activation and diverse functions: Motojima K, Cell Struct Funct. 1993 Oct; 18(5): 267-77).

Two variants of PPARgamma exist, PPARgamma₁ and gamma₂, which are the
5 result of alternative use of promoters and differential mRNA splicing (Vidal-Puig et al. J. Clin. Invest., 97:2553-2561, 1996). Different PPARs have different tissue distribution and modulate different physiological functions. The PPARs play a key role in various aspects of the regulation of a large number of genes, the products of which genes are directly or indirectly crucially involved in lipid and carbohydrate
10 metabolism. Thus, for example, PPARalpha receptors play an important part in the regulation of fatty acid catabolism or lipoprotein metabolism in the liver, while PPARgamma is crucially involved for example in regulating adipose cell differentiation. In addition, however, PPARs are also involved in the regulation of many other physiological processes, including those which are not directly
15 connected with carbohydrate or lipid metabolism. The activity of different PPARs can be modulated by various fatty acids, fatty acid derivatives and synthetic compounds to varying extents. For relevant reviews about functions, physiological effect and pathophysiology, see: Joel Berger et al., Annu. Rev. Med. 2002, 53, 409 – 435; Timothy Wilson et al. J. Med. Chem., 2000, Vol. 43, No. 4, 527-550; Steven
20 Kliewer et al., Recent Prog Horm Res. 2001; 56: 239-63.

The present invention relates to compounds of the formula I suitable for modulating the activity of PPARs, especially the activity of PPARalpha and PPARgamma. Depending on the modulation profile, the compounds of the formula
25 I are suitable for the treatment, control and prophylaxis of the indications described hereinafter, and for a number of other pharmaceutical applications connected thereto (see, for example, Joel Berger et al., Annu. Rev. Med. 2002, 53, 409 – 435; Timothy Wilson et al. J. Med. Chem., 2000, Vol. 43, No. 4, 527-550; Steven Kliewer et al., Recent Prog Horm Res. 2001; 56: 239-63; Jean-Charles Fruchart,
30 Bart Staels and Patrick Duriez: PPARs, Metabolic Disease and Arteriosclerosis, Pharmacological Research, Vol. 44, No. 5, 345-52; 2001; Sander Kersten, Beatrice Desvergne & Walter Wahli: Roles of PPARs in health and disease, NATURE, VOL 405, 25 MAY 2000; 421-4; Ines Pineda Torra, Giulia Chinetti,

Caroline Duval, Jean-Charles Fruchart and Bart Staels: Peroxisome proliferator-activated receptors: from transcriptional control to clinical practice, Curr Opin Lipidol 12: 2001, 245-254).

5 Compounds of this type are particularly suitable for the treatment and/or prevention of

1. - disorders of fatty acid metabolism and glucose utilization disorders
- disorders in which insulin resistance is involved

10

2. Diabetes mellitus, especially type 2 diabetes, including the prevention of the sequelae associated therewith.

Particular aspects in this connection are

- hyperglycemia,
- 15 - improvement in insulin resistance,
- improvement in glucose tolerance,
- protection of the pancreatic β cells
- prevention of macro- and microvascular disorders
3. Dyslipidemias and their sequelae such as, for example, atherosclerosis,
- 20 coronary heart disease, cerebrovascular disorders etc, especially those (but not restricted thereto) which are characterized by one or more of the following factors:
- high plasma triglyceride concentrations, high postprandial plasma triglyceride concentrations,
- 25 - low HDL cholesterol concentrations
- low ApoA lipoprotein concentrations
- high LDL cholesterol concentrations
- small dense LDL cholesterol particles
- high ApoB lipoprotein concentrations

30

4. Various other conditions which may be associated with the metabolic syndrome, such as:

- obesity (excess weight), including central obesity

- thromboses, hypercoagulable and prothrombotic states (arterial and venous)
- high blood pressure
- heart failure such as, for example (but not restricted thereto), following myocardial infarction, hypertensive heart disease or cardiomyopathy

5
10
15
20
25
30

5. Other disorders or conditions in which inflammatory reactions or cell differentiation may for example be involved are:

- atherosclerosis such as, for example (but not restricted thereto), coronary sclerosis including angina pectoris or myocardial infarction, stroke
- vascular restenosis or reocclusion
- chronic inflammatory bowel diseases such as, for example, Crohn's disease and ulcerative colitis
- pancreatitis
- other inflammatory states
- retinopathy
- adipose cell tumors
- lipomatous carcinomas such as, for example, liposarcomas
- solid tumors and neoplasms such as, for example (but not restricted thereto), carcinomas of the gastrointestinal tract, of the liver, of the biliary tract and of the pancreas, endocrine tumors, carcinomas of the lungs, of the kidneys and the urinary tract, of the genital tract, prostate carcinomas etc
- acute and chronic myeloproliferative disorders and lymphomas
- angiogenesis
- neurodegenerative disorders
- Alzheimer's disease
- multiple sclerosis
- Parkinson's disease
- erythematous-squamous dermatoses such as, for example, psoriasis
- acne vulgaris
- other skin disorders and dermatological conditions which are modulated by PPAR
- eczemas and neurodermitis

- dermatitis such as, for example, seborrheic dermatitis or photodermatitis
- keratitis and keratoses such as, for example, seborrheic keratoses, senile keratoses, actinic keratosis, photo-induced keratoses or keratosis follicularis
- 5 - keloids and keloid prophylaxis
- warts, including condylomata or condylomata acuminata
- human papilloma viral (HPV) infections such as, for example, venereal papillomata, viral warts such as, for example, molluscum contagiosum, leukoplakia
- 10 - papular dermatoses such as, for example, Lichen planus
- skin cancer such as, for example, basal-cell carcinomas, melanomas or cutaneous T-cell lymphomas
- localized benign epidermal tumors such as, for example, keratoderma, epidermal naevi
- 15 - chilblains
- high blood pressure
- syndrome X
- polycystic ovary syndrome (PCOS)
- asthma
- 20 - osteoarthritis
- lupus erythematosus (LE) or inflammatory rheumatic disorders such as, for example, rheumatoid arthritis
- vasculitis
- wasting (cachexia)
- 25 - gout
- ischemia/reperfusion syndrome
- acute respiratory distress syndrome (ARDS)

Formulations

30

The amount of a compound of formula I necessary to achieve the desired biological effect depends on a number of factors, for example the specific compound chosen, the intended use, the mode of administration and the clinical

condition of the patient. The daily dose is generally in the range from 0.001 mg to 100 mg (typically from 0.01 mg to 50 mg) per day and per kilogram of bodyweight, for example 0.1-10 mg/kg/day. An intravenous dose may be, for example, in the range from 0.001 mg to 1.0 mg/kg, which can suitably be administered as infusion

5 of 10 ng to 100 ng per kilogram and per minute. Suitable infusion solutions for these purposes may contain, for example, from 0.1 ng to 10 mg, typically from 1 ng to 10 mg, per milliliter. Single doses may contain, for example, from 1 mg to 10 g of the active ingredient. Thus, ampules for injections may contain, for example, from 1 mg to 100 mg, and single-dose formulations which can be

10 administered orally, such as, for example, capsules or tablets, may contain, for example, from 0.05 to 1000 mg, typically from 0.5 to 600 mg. For the therapy of the abovementioned conditions, the compounds of formula I may be used as the compound itself, but they are preferably in the form of a pharmaceutical composition with an acceptable carrier. The carrier must, of course, be acceptable

15 in the sense that it is compatible with the other ingredients of the composition and is not harmful for the patient's health. The carrier may be a solid or a liquid or both and is preferably formulated with the compound as a single dose, for example as a tablet, which may contain from 0.05% to 95% by weight of the active ingredient. Other pharmaceutically active substances may likewise be present, including other

20 compounds of formula I. The pharmaceutical compositions of the invention can be produced by one of the known pharmaceutical methods, which essentially consist of mixing the ingredients with pharmacologically acceptable carriers and/or excipients.

25 Pharmaceutical compositions of the invention are those suitable for oral, rectal, topical, peroral (for example sublingual) and parenteral (for example subcutaneous, intramuscular, intradermal or intravenous) administration, although the most suitable mode of administration depends in each individual case on the nature and severity of the condition to be treated and on the nature of the

30 compound of formula I used in each case. Coated formulations and coated slow-release formulations also belong within the framework of the invention. Preference is given to acid- and gastric juice-resistant formulations. Suitable coatings resistant to gastric juice comprise cellulose acetate phthalate, polyvinyl acetate phthalate,

hydroxypropylmethylcellulose phthalate and anionic polymers of methacrylic acid and methyl methacrylate.

Suitable pharmaceutical preparations for oral administration may be in the form of
5 separate units such as, for example, capsules, cachets, suckable tablets or
tablets, each of which contain a defined amount of the compound of formula I; as
powders or granules, as solution or suspension in an aqueous or nonaqueous
liquid; or as an oil-in-water or water-in-oil emulsion. These compositions may, as
10 already mentioned, be prepared by any suitable pharmaceutical method which
includes a step in which the active ingredient and the carrier (which may consist of
one or more additional ingredients) are brought into contact. The compositions are
generally produced by uniform and homogeneous mixing of the active ingredient
with a liquid and/or finely divided solid carrier, after which the product is shaped if
15 necessary. Thus, for example, a tablet can be produced by compressing or
molding a powder or granules of the compound, where appropriate with one or
more additional ingredients. Compressed tablets can be produced by tableting the
compound in free-flowing form such as, for example, a powder or granules, where
appropriate mixed with a binder, glidant, inert diluent and/or one (or more) surface-
20 active/dispersing agent(s) in a suitable machine. Molded tablets can be produced
by molding the compound, which is in powder form and is moistened with an inert
liquid diluent, in a suitable machine.

Pharmaceutical compositions which are suitable for peroral (sublingual)
administration comprise suckable tablets which contain a compound of formula I
25 with a flavoring, normally sucrose and gum arabic or tragacanth, and pastilles
which comprise the compound in an inert base such as gelatin and glycerol or
sucrose and gum arabic.

Pharmaceutical compositions suitable for parenteral administration comprise
30 preferably sterile aqueous preparations of a compound of formula I, which are
preferably isotonic with the blood of the intended recipient. These preparations are
preferably administered intravenously, although administration may also take
place by subcutaneous, intramuscular or intradermal injection. These preparations

can preferably be produced by mixing the compound with water and making the resulting solution sterile and isotonic with blood. Injectable compositions of the invention generally contain from 0.1 to 5% by weight of the active compound.

- 5 Pharmaceutical compositions suitable for rectal administration are preferably in the form of single-dose suppositories. These can be produced by mixing a compound of the formula I with one or more conventional solid carriers, for example cocoa butter, and shaping the resulting mixture.
- 10 Pharmaceutical compositions suitable for topical use on the skin are preferably in the form of ointment, cream, lotion, paste, spray, aerosol or oil. Carriers which can be used are petrolatum, lanolin, polyethylene glycols, alcohols and combinations of two or more of these substances. The active ingredient is generally present in a concentration of from 0.1 to 15% by weight of the composition, for example from
- 15 0.5 to 2%.

- Transdermal administration is also possible. Pharmaceutical compositions suitable for transdermal uses can be in the form of single plasters which are suitable for long-term close contact with the patient's epidermis. Such plasters suitably contain
- 20 the active ingredient in an aqueous solution which is buffered where appropriate, dissolved and/or dispersed in an adhesive or dispersed in a polymer. A suitable active ingredient concentration is about 1% to 35%, preferably about 3% to 15%. A particular possibility is for the active ingredient to be released by electrotransport or iontophoresis as described, for example, in Pharmaceutical Research, 2(6): 318
- 25 (1986).

- The compounds of the formula I are distinguished by favorable effects on metabolic disorders. They beneficially influence lipid and sugar metabolism, in particular they lower the triglyceride level and are suitable for the prevention and
- 30 treatment of type II diabetes and arteriosclerosis and the diverse sequelae thereof.

Combinations with other medicaments

The compounds of the invention can be administered alone or in combination with one or more further pharmacologically active substances which have, for example, favorable effects on metabolic disturbances or disorders frequently associated therewith. Examples of such medicaments are

1. medicaments which lower blood glucose, antidiabetics,
2. active ingredients for the treatment of dyslipidemias,
3. antiatherosclerotic medicaments,
4. antiobesity agents,
5. antiinflammatory active ingredients
6. active ingredients for the treatment of malignant tumors
7. antithrombotic active ingredients
8. active ingredients for the treatment of high blood pressure
9. active ingredients for the treatment of heart failure and active ingredients for the treatment and/or prevention of complications caused by diabetes or associated with diabetes.

They can be combined with the compounds of the invention of the formula I in particular for a synergistic improvement in the effect. Administration of the active ingredient combination can take place either by separate administration of the active ingredients to the patient or in the form of combination products in which a plurality of active ingredients are present in one pharmaceutical preparation.

Examples which may be mentioned are:

Antidiabetics


Suitable antidiabetics are disclosed for example in the Rote Liste 2001, chapter 12 or in the USP Dictionary of USAN and International Drug Names, US Pharmacopeia, Rockville 2001. Antidiabetics include all insulins and insulin derivatives such as, for example, Lantus[®] (see www.lantus.com) or Apidra[®], and other fast-acting insulins (see US 6,221,633), GLP-1 receptor modulators as

described in WO 01/04146 or else, for example, those disclosed in WO 98/08871 of Novo Nordisk A/S.

The orally effective hypoglycemic active ingredients include, preferably,
5 sulfonyleureas, biguanides, meglitinides, oxadiazolidinediones, thiazolidinediones, glucosidase inhibitors, glucagon antagonists, GLP-1 agonists, DPP-IV inhibitors, potassium channel openers such as, for example, those disclosed in
WO 97/26265 and WO 99/03861, insulin sensitizers, inhibitors of liver enzymes
involved in the stimulation of gluconeogenesis and/or glycogenolysis, modulators
10 of glucose uptake, compounds which alter lipid metabolism and lead to a change in the blood lipid composition, compounds which reduce food intake, PPAR and PXR modulators and active ingredients which act on the ATP-dependent potassium channel of the beta cells.

15 In one embodiment of the invention, the compounds of the formula I are administered in combination with insulin.

In one embodiment of the invention, the compounds of the formula I are administered in combination with substances which influence hepatic glucose
20 production such as, for example, glycogen phosphorylase inhibitors (see: WO 01/94300, WO 02/096864, WO 03/084923, WO 03/084922, WO 03/104188)

 In one embodiment, the compounds of the formula I are administered in combination with a sulfonyleurea such as, for example, tolbutamide, glibenclamide,
25 glipizide or glimepiride.

In one embodiment, the compounds of the formula I are administered in combination with an active ingredient which acts on the ATP-dependent potassium channel of the beta cells, such as, for example, tolbutamide, glibenclamide,
30 glipizide, glimepiride or repaglinide.

In one embodiment, the compounds of the formula I are administered in combination with a biguanide such as, for example, metformin.

In a further embodiment, the compounds of the formula I are administered in combination with a meglitinide such as, for example, repaglinide.

5 In one embodiment, the compounds of the formula I are administered in combination with a thiazolidinedione such as, for example, ciglitazone, pioglitazone, rosiglitazone or the compounds disclosed in WO 97/41097 of Dr. Reddy's Research Foundation, in particular 5-[[4-[(3,4-dihydro-3-methyl-4-oxo-2-quinazolinylmethoxy)phenyl]methyl]-2,4-thiazolidinedione.

10 In one embodiment, the compounds of the formula I are administered in combination with a DPPIV inhibitor as described, for example, in WO98/19998, WO99/61431, WO99/67278, WO99/67279, WO01/72290, WO 02/38541, WO03/040174, in particular P 93/01 (1-cyclopentyl-3-methyl-1-oxo-2-pentan ammonium chloride), P-31/98, LAF237 (1-[2-[3-hydroxyadamant-1-ylamino)acetyl]pyrrolidine-2-(S)-carbonitrile), TS021 ((2S, 4S)-4-fluoro-1-[(2-hydroxy-1,1-dimethylethyl)amino]-acetyl]pyrrolidine-2-carbonitrile monobenzenesulfonate).

15 In one embodiment of the invention, the compounds of the formula I are administered in combination with a PPARgamma agonist such as, for example, rosiglitazone, pioglitazone.

20 In one embodiment, the compounds of the formula I are administered in combination with compounds with an inhibitory effect on SGLT-1 and/or 2, as disclosed directly or indirectly for example in PCT/EP03/06841, PCT/EP03/13454 and PCT/EP03/13455.

25 In one embodiment, the compounds of the formula I are administered in combination with an α -glucosidase inhibitor such as, for example, miglitol or acarbose.

30 In one embodiment, the compounds of the formula I are administered in combination with more than one of the aforementioned compounds, e.g. in

combination with a sulfonylurea and metformin, a sulfonylurea and acarbose, repaglinide and metformin, insulin and a sulfonylurea, insulin and metformin, insulin and troglitazone, insulin and lovastatin, etc.

5 Lipid modulators

In one embodiment of the invention, the compounds of the formula I are administered in combination with an HMGCoA reductase inhibitor such as lovastatin, fluvastatin, pravastatin, simvastatin, ivastatin, itavastatin, atorvastatin,
10 rosuvastatin.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a bile acid reabsorption inhibitor (see, for example, US 6,245,744, US 6,221,897, US 6,277,831, EP 0683 773, EP 0683
15 774).

In one embodiment of the invention, the compounds of the formula I are administered in combination with a polymeric bile acid adsorbent such as, for example, cholestyramine, colessevelam.
20

In one embodiment of the invention, the compounds of the formula I are administered in combination with a cholesterol absorption inhibitor as described for example in WO 0250027, or ezetimibe, tiqueside, pamaqueside.

25 In one embodiment of the invention, the compounds of the formula I are administered in combination with an LDL receptor inducer (see, for example, US 6,342,512).

In one embodiment, the compounds of the formula I are administered in
30 combination with bulking agents, preferably insoluble bulking agents (see, for example, carob/Caromax® (Zunft H J; et al., Carob pulp preparation for treatment of hypercholesterolemia, ADVANCES IN THERAPY (2001 Sep-Oct), 18(5), 230-6.) Caromax is a carob-containing product from Nutrinova, Nutrition Specialties &

Food Ingredients GmbH, Industriepark Höchst, 65926 Frankfurt/Main)).

Combination with Caromax® is possible in one preparation or by separate administration of compounds of the formula I and Caromax®. Caromax® can in this connection also be administered in the form of food products such as, for example, in bakery products or muesli bars.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a PPARalpha agonist.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a mixed PPAR alpha/gamma agonist such as, for example, AZ 242 (Tesaglitazar, (S)-3-(4-[2-(4-methanesulfonyloxyphenyl)ethoxy]phenyl)-2-ethoxypropionic acid), BMS 298585 (N-[(4-methoxyphenoxy)carbonyl]-N-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]phenyl]methylglycine) or as described in WO 99/62872, WO 99/62871, WO 01/40171, WO 01/40169, WO96/38428, WO 01/81327, WO 01/21602, WO 03/020269, WO 00/64888 or WO 00/64876.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a fibrate such as, for example, fenofibrate, gemfibrozil, clofibrate, bezafibrate.

In one embodiment of the invention, the compounds of the formula I are administered in combination with nicotinic acid or niacin.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a CETP inhibitor, e.g. CP- 529, 414 (torcetrapib).

In one embodiment of the invention, the compounds of the formula I are administered in combination with an ACAT inhibitor.

In one embodiment of the invention, the compounds of the formula I are administered in combination with an MTP inhibitor such as, for example, implitapide.

In one embodiment of the invention, the compounds of the formula I are administered in combination with an antioxidant.

- 5 In one embodiment of the invention, the compounds of the formula I are administered in combination with a lipoprotein lipase inhibitor.

In one embodiment of the invention, the compounds of the formula I are administered in combination with an ATP citrate lyase inhibitor.

10

In one embodiment of the invention, the compounds of the formula I are administered in combination with a squalene synthetase inhibitor.

- 15 In one embodiment of the invention, the compounds of the formula I are administered in combination with a lipoprotein(a) antagonist.

Antiobesity agents

- 20 In one embodiment of the invention, the compounds of the formula I are administered in combination with a lipase inhibitor such as, for example, orlistat.

In one embodiment, the further active ingredient is fenfluramine or dexfenfluramine.

- 25 In another embodiment, the further active ingredient is sibutramine.

- In a further embodiment, the compounds of the formula I are administered in combination with CART modulators (see "Cocaine-amphetamine-regulated transcript influences energy metabolism, anxiety and gastric emptying in mice" Asakawa, A, et al., M.: Hormone and Metabolic Research (2001), 33(9), 554-558). NPY antagonists, e.g. naphthalene-1-sulfonic acid {4-[(4-aminoquinazolin-2-ylamino)methyl]- cyclohexylmethyl}amide hydrochloride (CGP 71683A)), MC4 agonists (e.g. 1-amino-1,2,3,4-tetrahydronaphthalene-2-carboxylic acid [2-(3a-
- 30

- benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydropyrazolo[4,3-c]pyridin-5-yl)-1-(4-chlorophenyl)-2-oxoethyl]-amide; (WO 01/91752)), orexin antagonists (e.g. 1-(2-methylbenzoxazol-6-yl)-3-[1,5]naphthyridin-4-ylurea hydrochloride (SB-334867-A)), H3 agonists (3-cyclohexyl-1-(4,4-dimethyl-1,4,6,7-tetrahydroimidazo[4,5-c]pyridin-5-yl)propan-1-one oxalic acid salt (WO 00/63208)); TNF agonists, CRF antagonists (e.g. [2-methyl-9-(2,4,6-trimethylphenyl)-9H-1,3,9-triazafluoren-4-yl]dipropylamine (WO 00/66585)), CRF BP antagonists (e.g. urocortin), urocortin agonists, $\beta 3$ agonists (e.g. 1-(4-chloro-3-methanesulfonylmethylphenyl)-2-[2-(2,3-dimethyl-1H-indol-6-yloxy)ethylamino]-ethanol hydrochloride (WO 01/83451)),
- 10 MSH (melanocyte-stimulating hormone) agonists, CCK-A agonists (e.g. {2-[4-(4-chloro-2,5-dimethoxyphenyl)-5-(2-cyclohexylethyl)thiazol-2-ylcarbamoyl]-5,7-dimethylindol-1-yl}acetic acid trifluoroacetic acid salt (WO 99/15525)), serotonin reuptake inhibitors (e.g. dexfenfluramine), mixed serotoninergic and noradrenergic compounds (e.g. WO 00/71549), 5HT agonists e.g. 1-(3-ethylbenzofuran-7-
- 15 yl)piperazine oxalic acid salt (WO 01/09111), bombesin agonists, galanin antagonists, growth hormone (e.g. human growth hormone), growth hormone-releasing compounds (6-benzoyloxy-1-(2-diisopropylaminoethylcarbonyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tertiary butyl ester (WO 01/85695)), TRH agonists (see, for example, EP 0 462 884), uncoupling protein 2 or 3 modulators,
- 20 leptin agonists (see, for example, Lee, Daniel W.; Leinung, Matthew C.; Rozhavskaya-Arena, Marina; Grasso, Patricia. Leptin agonists as a potential approach to the treatment of obesity. *Drugs of the Future* (2001), 26(9), 873-881), DA agonists (bromocriptine, Doprexin), lipase/amylase inhibitors (e.g. WO 00/40569), PPAR modulators (e.g. WO 00/78312), RXR modulators or TR- β
- 25 agonists.

In one embodiment of the invention, the further active ingredient is leptin.

- In one embodiment, the further active ingredient is dexamphetamine,
- 30 amphetamine, mazindole or phentermine.

In one embodiment, the compounds of the formula I are administered in combination with medicaments having effects on the coronary circulation and the

vascular system, such as, for example, ACE inhibitors (e.g. ramipril), medicaments which act on the angiotensin-renaline system, calcium antagonists, beta blockers etc.

- 5 In one embodiment, the compounds of the formula I are administered in combination with medicaments having an antiinflammatory effect.

In one embodiment, the compounds of the formula I are administered in combination with medicaments which are employed for cancer therapy and cancer
10 prevention.

It will be appreciated that every suitable combination of the compounds of the invention with one or more of the aforementioned compounds and optionally one or more other pharmacologically active substances is regarded as falling within the
15 protection conferred by the present invention.

The activity of the compounds was tested as follows:

20 **Determination of EC50 values of PPAR agonists in the cellular PPARalpha assay**

Principle

The potency of substances which bind to human PPARalpha and activate in an
25 agonistic manner is analyzed using a stably transfected HEK cell line (HEK=human emryo kidney) which is referred to here as PPARalpha reporter cell line. It contains two genetic elements, a luciferase reporter element (pdeltaM-GAL4-Luc-Zeo) and a PPARalpha fusion protein (GR-GAL4-humanPPARalpha-LBD) which mediates expression of the luciferase reporter element depending on a
30 PPARalpha ligand. The stably and constitutively expressed fusion protein GR-GAL4-humanPPARalpha-LBD binds in the cell nucleus of the PPARalpha reporter cell line via the GAL4 protein portion to the GAL4 DNA binding motifs 5'-upstream of the luciferase reporter element which is integrated in the genome of the cell line.

There is only little expression of the luciferase reporter gene without addition of a PPARalpha ligand if fatty acid-depleted fetal calf serum (cs-FCS) is used in the assay. PPARalpha ligands bind and activate the PPARalpha fusion protein and thereby bring about expression of the luciferase reporter gene. The luciferase
5 which is formed can be detected by means of chemiluminescence via an appropriate substrate.

Construction of the cell line

10 The PPARalpha reporter cell line was prepared in 2 stages. Firstly, the luciferase reporter element was constructed and stably transfected into HEK cells. For this purpose, five binding sites of the yeast transcription factor GAL4 (each 5'-CGGAGTACTGTCTCCGAG-3') were cloned in 5'-upstream of a 68 bp-long minimal MMTV promoter (Genbank Accession # V01175). The minimal MMTV
15 promoter section contains a CCAAT box and a TATA element in order to enable efficient transcription by RNA polymerase II. The cloning and sequencing of the GAL4-MMTV construct took place in analogy to the description of Sambrook J. *et al.* (Molecular cloning, Cold Spring Harbor Laboratory Press, 1989). Then the complete Photinus pyralis gene (Genbank Accession # M15077) was cloned in 3'-
20 downstream of the GAL4-MMTV element. After sequencing, the luciferase reporter element consisting of five GAL4 binding sites, MMTV promoter and luciferase gene was recloned into a plasmid which confers zeocin resistance in order to obtain the plasmid pdeltaM-GAL4-Luc-Zeo. This vector was transfected into HEK cells in accordance with the statements in Ausubel, F.M. *et al.* (Current
25 protocols in molecular biology, Vol. 1-3, John Wiley & Sons, Inc., 1995). Then zeocin-containing medium (0.5 mg/ml) was used to select a suitable stable cell clone which showed very low basal expression of the luciferase gene.

In a second step, the PPARalpha fusion protein (GR-GAL4-humanPPARalpha-LBD) was introduced into the stable cell clone described. For this purpose, initially
30 the cDNA coding for the N-terminal 76 amino acids of the glucocorticoid receptor (Genbank Accession # P04150) was linked to the cDNA section coding for amino acids 1-147 of the yeast transcription factor GAL4 (Genbank Accession # P04386). The cDNA of the ligand-binding domain of the human PPARalpha

receptor (amino acids S167-Y468; Genbank Accession # S74349) was cloned in at the 3'-end of this GR-GAL4 construct. The fusion construct prepared in this way (GR-GAL4-humanPPARalpha-LBD) was recloned into the plasmid pcDNA3 (from Invitrogen) in order to enable constitutive expression therein by the
5 cytomagalovirus promoter. This plasmid was linearized with a restriction endonuclease and stably transfected into the previously described cell clone containing the luciferase reporter element. The finished PPARalpha reporter cell line which contains a luciferase reporter element and constitutively expresses the PPARalpha fusion protein (GR-GAL4-human PPARalpha-LBD) was isolated by
10 selection with zeocin (0.5 mg/ml) and G418 (0.5 mg/ml).

Assay procedure

The activity of PPARalpha agonists is determined in a 3-day assay which is
15 described below:

Day 1

The PPARalpha reporter cell line is cultivated to 80% confluence in DMEM (# 41965-039, Invitrogen) which is mixed with the following additions: 10% cs-FCS
20 (fetal calf serum; #SH-30068.03, Hyclone), 0.5 mg/ml zeocin (#R250-01, Invitrogen), 0.5 mg/ml G418 (#10131-027, Invitrogen), 1% penicillin-streptomycin solution (#15140-122, Invitrogen) and 2 mM L-glutamine (#25030-024, Invitrogen). The cultivation takes place in standard cell culture bottles (# 353112, Becton Dickinson) in a cell culture incubator at 37°C in the presence of 5% CO₂. The
25 80%-confluent cells are washed once with 15 ml of PBS (#14190-094, Invitrogen), treated with 3 ml of trypsin solution (#25300-054, Invitrogen) at 37°C for 2 min, taken up in 5 ml of the DMEM described and counted in a cell counter. After dilution to 500,000 cells/ml, 35,000 cells are seeded in each well of a 96 well microtiter plate with a clear plastic base (#3610, Corning Costar). The plates are
30 incubated in the cell culture incubator at 37°C and 5% CO₂ for 24 h.

Day 2

PPARalpha agonists to be tested are dissolved in DMSO in a concentration of

10 mM. This stock solution is diluted in DMEM (#41965-039, Invitrogen) which is mixed with 5% cs-FCS (#SH-30068.03, Hyclone), 2 mM L-glutamine (#25030-024, Invitrogen) and the previously described antibiotics (zeocin, G418, penicillin and streptomycin).

- 5 Test substances are tested in 11 different concentrations in the range from 10 μ M to 100 pM. More potent compounds are tested in concentration ranges from 1 μ M to 10 pM or between 100 nM and 1 pM.

The medium of the PPARAlpha reporter cell line seeded on day 1 is completely removed by aspiration, and the test substances diluted in medium are immediately
10 added to the cells. The dilution and addition of the substances is carried out by a robot (Beckman FX). The final volume of the test substances diluted in medium is 100 μ l per well of a 96 well microtiter plate. The DMSO concentration in the assay is less than 0.1 % v/v in order to avoid cytotoxic effects of the solvent.

Each plate was charged with a standard PPARAlpha agonist, which was likewise
15 diluted in 11 different concentrations, in order to demonstrate the functioning of the assay in each individual plate. The assay plates are incubated in an incubator at 37°C and 5% CO₂ for 24 h.

Day 3

- 20 The PPARAlpha reporter cells treated with the test substances are removed from the incubator, and the medium is aspirated off. The cells are lyzed by pipetting 50 μ l of Bright Glo reagent (from Promega) into each well of a 96 well microtiter plate. After incubation at room temperature in the dark for 10 minutes, the microtiter plates are measured in the luminometer (Trilux from Wallac). The measuring time
25 for each well of a microtiter plate is 1 sec.

Evaluation

- The raw data from the luminometer are transferred into a Microsoft Excel file.
30 Dose-effect plots and EC50 values of PPAR agonists are calculated using the XL.Fit program as specified by the manufacturer (IDBS).

The PPARAlpha EC50 values for the compounds of Examples 1 to 13 in this assay

are in the range from 0.05nM to >10 μ M.

The results for the activity of some compounds of the invention of the formula I are indicated in Table I below:

5

Table I

Example No.	EC50 PPARalpha [nM]
1	0.2
3	0.2
4	0.6
10	0.3
11	27
12	34
13	0.06

10 It is evident from Table I that the compounds of the invention of the formula I activate the PPARalpha receptor and thus bring about for example in analogy to fibrates in clinical use a lowering of triglycerides in the body (see, for example, J.-Ch. Fruchard et al.: PPARs, Metabolic Disease and Atherosclerosis, Pharmacological Research, Vol. 44, No. 5, 345-52, 2001; S. Kersten et al.: Roles of PPARs in health and disease, NATURE, VOL 405, 25 MAY 2000, 421-4; I. Pineda et al.: Peroxisome proliferator-activated receptors: from transcriptional control to clinical practice, Curr Opin Lipidol 12: 2001, 245-254).

15 Determination of EC50 values of PPAR agonists in the cellular PPARgamma assay

20

Principle

A transient transfection system is employed to determine the cellular PPARgamma activity of PPAR agonists. It is based on the use of a luciferase reporter plasmid (pGL3basic-5xGAL4-TK) and of a PPARgamma expression plasmid (pcDNA3-GAL4-humanPPARgammaLBD). Both plasmids are transiently transfected into human embryonic kidney cells (HEK cells). There is then expression in these cells of the fusion protein GAL4-humanPPARgammaLBD which binds to the GAL4 binding sites of the reporter plasmid. In the presence of a PPARgamma-active ligand, the activated fusion protein GAL4-humanPPARgammaLBD induces expression of the luciferase reporter gene, which can be detected in the form of a chemiluminescence signal after addition of a luciferase substrate. As a difference from the stably transfected PPARalpha reporter cell line, in the cellular PPARgamma assay the two components (luciferase reporter plasmid and PPARgamma expression plasmid) are transiently transfected into HEK cells because stable and permanent expression of the PPARgamma fusion protein is cytotoxic.

Construction of the plasmids

The luciferase reporter plasmid pGL3basic-5xGAL4-TK is based on the vector pGL3basic from Promega. The reporter plasmid is prepared by cloning five binding sites of the yeast transcription factor GAL4 (each binding site with the sequence 5'-CTCGGAGGACAGTACTCCG-3'), together with a 160 bp-long thymidine kinase promoter section (Genbank Accession # AF027128) 5'-upstream into pGL3basic. 3'-downstream of the thymidine kinase promoter is the complete luciferase gene from *Photinus pyralis* (Genbank Accession # M15077) which is already a constituent of the plasmid pGL3basic used. The cloning and sequencing of the reporter plasmid pGL3basic-5xGAL4-TK took place in analogy to the description in Sambrook J. *et. al.* (Molecular cloning, Cold Spring Harbor Laboratory Press, 1989).

The PPARgamma expression plasmid pcDNA3-GAL4-humanPPARgammaLBD was prepared by first cloning the cDNA coding for amino acids 1-147 of the yeast transcription factor GAL4 (Genbank Accession # P04386) into the plasmid pcDNA3 (from Invitrogen) 3'-downstream of the cytomegalovirus promoter.

Subsequently, the cDNA of the ligand-binding domain (LBD) of the human PPARgamma receptor (amino acids 1152-Y475; Accession # g1480099) 3'-downstream of the GAL4 DNA binding domain. Cloning and sequencing of the PPARgamma expression plasmid pcDNA3-GAL4-humanPPARgammaLBD again
5 took place in analogy to the description in Sambrook J. *et. al.* (Molecular cloning, Cold Spring Harbor Laboratory Press, 1989). Besides the luciferase reporter plasmid pGL3basic-5xGAL4-TK and the PPARgamma expression plasmid pcDNA3-GAL4-humanPPARgammaLBD, also used for the cellular PPARgamma assay are the reference plasmid pRL-CMV (from Promega) and the plasmid
10 pBluescript SK(+) from Stratagene. All four plasmids were prepared using a plasmid preparation kit from Qiagen, which ensured a plasmid quality with a minimal endotoxin content, before transfection into HEK cells.

15 Assay procedure

The activity of PPARgamma agonists is determined in a 4-day assay which is described below. Before the transfection, HEK cells are cultivated in DMEM (# 41965-039, Invitrogen) which is mixed with the following additions: 10% FCS (#16000-044, Invitrogen), 1% penicillin-streptomycin solution (#15140-122,
20 Invitrogen) and 2 mM L-glutamine (#25030-024, Invitrogen).

Day 1

Firstly, solution A, a transfection mixture which contains all four plasmids previously described in addition to DMEM, is prepared. The following amounts are
25 used to make up 3 ml of solution A for each 96 well microtiter plate for an assay: 2622 µl of antibiotic- and serum-free DMEM (# 41965-039, Invitrogen), 100 µl of reference plasmid pRL-CMV (1 ng/µl), 100 µl of luciferase reporter plasmid pGL3basic-5xGAL4-TK (10 ng/µl), 100 µl of PPARgamma expression plasmid pcDNA3-GAL4-humanPPARgammaLBD (100 ng/µl) and 78 µl of plasmid
30 pBluescript SK(+) (500 ng/µl). Then 2 ml of solution B are prepared by mixing 1.9 ml of DMEM (# 41965-039, Invitrogen) with 100 µl of PolyFect transfection reagent (from Qiagen) for each 96 well microtiter plate. Subsequently, 3 ml of solution A are mixed with 2 ml of solution B to give 5 ml of solution C, which is thoroughly

mixed by multiple pipetting and incubated at room temperature for 10 min.

80%-confluent HEK cells from a cell culture bottle with a capacity of 175 cm² are washed once with 15 ml of PBS (#14190-094, Invitrogen) and treated with 3 ml of trypsin solution (#25300-054, Invitrogen) at 37°C for 2 min. The cells are then
5 taken up in 15 ml of DMEM (# 41965-039, Invitrogen) which is mixed with 10% FCS (# 16000-044, Invitrogen), 1% penicillin-streptomycin solution (#15140-122, Invitrogen) and 2 mM L-glutamine (#25030-024, Invitrogen). After the cell suspension has been counted in a cell counter, the suspension is diluted to
10 250,000 cells/ml. 15 ml of this cell suspension are mixed with 5 ml of solution C for one microtiter plate. 200 µl of the suspension are seeded in each well of a 96 well microtiter plate with a clear plastic base (#3610, Corning Costar). The plates are incubated in a cell culture incubator at 37°C and 5% CO₂ for 24 h.

Day 2

15 PPAR agonists to be tested are dissolved in DMSO in a concentration of 10 mM. This stock solution is diluted in DMEM (# 41965-039, Invitrogen) which is mixed with 2% Ultrosor (#12039-012, Biosepra), 1% penicillin-streptomycin solution (#15140-122, Invitrogen) and 2 mM L-glutamine (#25030-024, Invitrogen). Test substances are tested in a total of 11 different concentrations in the range from 10
20 µM to 100 pM. More potent compounds are tested in concentration ranges from 1 µM to 10 pM.

The medium of the HEK cells transfected and seeded on day 1 is completely removed by aspiration, and the test substances diluted in medium are immediately added to the cells. The dilution and addition of the substances is carried out by a
25 robot (Beckman FX). The final volume of the test substances diluted in medium is 100 µl per well of a 96 well microtiter plate. Each plate is charged with a standard PPARgamma agonist, which is likewise diluted in 11 different concentrations, in order to demonstrate the functioning of the assay in each individual plate. The assay plates are incubated in an incubator at 37°C and 5% CO₂.

30

Day 4

After removal of the medium by aspiration, 50 µl of Dual-Glo™ reagent (Dual-Glo™ Luciferase Assay System; Promega) are added to each well in accordance

with the manufacturer's instructions in order to lyse the cells and provide the substrate for the firefly luciferase (*Photinus pyralis*) formed in the cells. After incubation at room temperature in the dark for 10 minutes, the firefly luciferase-mediated chemiluminescence is measured in a measuring instrument (measuring time/well 1 sec; Trilux from Wallac). Then 50 μ l of the Dual-Glo™ Stop & Glo reagent (Dual-Glo™ Luciferase Assay System; Promega) is added to each well in order to stop the activity of the firefly luciferase and provide the substrate for the Renilla luciferase expressed by the reference plasmid pRL-CMV. After incubation at room temperature in the dark for a further 10 minutes, a chemiluminescence mediated by the Renilla luciferase is again measured for 1 sec/well in the measuring instrument.


Evaluation

The crude data from the luminometer are transferred into a Microsoft Excel file. The firefly/Renilla luciferase activity ratio is determined for each measurement derived from one well of the microtiter plate. The dose-effect plots and EC50 values of PPAR agonists are calculated from the ratios by the XL.Fit program as specified by the manufacturer (IDBS).

PPARgamma EC50 values in the range from 0.5nM to >10 μ M were measured for the PPAR agonists described in this application.





The examples given below serve to illustrate the invention, but without limiting it.

33




Ex.	R1	R2	Ring B	R4	X	Ring A	Y	R3	R5
I	H	H		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
II	H	H		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
III	H	H		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me

34

Ex.	R1	R2	Ring B	R4	X	Ring A	Y	R3	R5
IV	5-Me	H		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
V	5-Me	H		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
VI	4-SCF3	H	Ph	Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
VII	3-OCF2-CF2H	H	Ph	Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
VIII	4-OPh	H	Ph	Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
IX	H	H		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
X	3-O-C2H4-O-Me	5-CF3		Me	CH2O	cis 1,3 Cy	CH2O	H	6-Me
XI	4-Me	H	Ph	Ph	CH2O	cis 1,3 Cy	CH2O	H	6-Me


35


Ex.	R1	R2	Ring B	R4	X	Ring A	Y	R3	R5
XII	3-OMe	H	Ph	Ph	CH ₂ O	cis 1,3 Cy	CH ₂ O	H	6-Me
XIII	H	H		H	CH ₂ O	cis 1,3 Cy	CH ₂ O	H	6-Me

cis 1,3 cy means: cis-substituted cyclohexane-1,3-diyl with Cahn-Ingold-Prelog stereochemistry as specified in the examples

----: indicates the point of attachment

The compounds of the formulae I and Ia can be obtained in accordance with the reaction scheme below:





Compounds of the formula A in which R3, R5 and Y have the meanings given above are reacted with NBS in an inert solvent (e.g. CCl₄), giving a compound of the formula B.

The compound of the formula B is reacted with a compound of the formula C in which n and m are each 0-5, giving a compound of the formula D in which R1, R2, R4, m, n and Y have the meanings described above; component C is initially heated with dibutyltin oxide in toluene on a water separator for a number of hours and then, with addition of dimethylformamide, cesium fluoride and bromide B, converted into D by stirring at room temperature for a number of hours.

The compound of the formula E is reacted with an aldehyde of the formula W (for example benzaldehyde, thiophene- or furancarbaldehyde) to give a compound of the formula F in which R1, R2, R4 and X are as defined above; to this end, components E and F are initially dissolved in acetic acid and HCl is introduced until the reaction has gone to completion, giving compounds of the formula F.

The compound of the formula F in which R1, R2, R4 and X are as defined above is heated under reflux with POCl₃ in chloroform for a number of hours, giving compounds of the formula G.

Compounds of the formula G in which R1, R2, R4 and X are as defined above are reacted with NaI in acetone under reflux for a number of hours, giving a compound of the formula H.

5

The compound of the formula D is reacted with a compound of formula H in which Y is as defined above, giving a compound of the formula J in which R1, R2, R4, R5, X and Y are as defined above. To establish an ether bond, D is deprotonated, for example, in a mixture of dimethylformamide and tetrahydrofuran using a strong base such as Na hydride, at room temperature, and then alkylated with component H.

10

The compound of the formula J is converted into compounds of the formula M in which R1, R2, R4, R5, X and Y are as defined above by hydrolyzing the ester function, for example by heating with potassium hydroxide in an alcohol (ethanol, tert-butanol), and releasing the carboxylic acid group of the formula I by acidification. This carboxylic acid group can be derivatized by customary methods into the group of the formula $-(C=O)-OR_3$ in which R3 is as defined above.

15


Other compounds can be obtained accordingly or by known processes.

20

Example

25

4,5-Dimethyl-2-naphthalen-2-ylloxazole 3-oxide




18.4 g of diacetyl monoxime and 31.2 g of 2-naphthaldehyde are added to 10 ml of

BEST AVAILABLE COPY


- glacial acetic acid, and HCl gas is introduced with ice-cooling for 30 minutes. The product is precipitated as the hydrochloride by addition of methyl tert-butyl ether and filtered off with suction, and the precipitate is washed with methyl tert-butyl ether. The precipitate is suspended in a mixture of dichloromethane and water, and a basic pH is established using ammonia. The mixture is extracted three times with in each case 500 ml of dichloromethane and ethyl acetate, the combined organic phases are dried over MgSO₄ and the solvent is then removed under reduced pressure. This gives 40.3 g of 4,5-dimethyl-2-naphthalen-2-yloxazole 3-oxide as a yellow solid.
- 10 CF15H13NO₂ (239.28), MS(ESI) = 240 (M+H⁺).

4-Chloromethyl-5-methyl-2-naphthalen-2-yloxazole



- 15 40 g of 4,5-dimethyl-2-naphthalen-2-yloxazole 3-oxide are dissolved in 200 ml of chloroform, 16.7 ml of phosphorus oxychloride are added and the mixture is heated under reflux for 30 minutes. The reaction mixture is cooled to 0°C, a slightly alkaline pH is established using ammonia and the mixture is extracted three times with in each case 500 ml of ethyl acetate. The combined organic phases are washed with water and dried over MgSO₄ and the solvent is then removed under reduced pressure. The residue is purified on silica gel using the mobile phase n-heptane:ethyl acetate = 80:1 => 5:1. This gives 10.6 g of 4-chloromethyl-5-methyl-2-naphthalen-2-yloxazole as a colorless solid. C₁₅H₁₂ClNO (257.72), MS(ESI):
- 20 258 (M+H⁺).
- 25


4-Iodomethyl-5-methyl-2-naphthalen-2-yloxazole



1.8 g of 4-chloromethyl-5-methyl-2-naphthalen-2-ylloxazole and 3 g of sodium iodide in 150 ml of acetone are heated under reflux for 2 hours. After the reaction mixture has been cooled, 300 ml of methyl tert-butyl ether are added, the mixture is washed three times with saturated Na₂S₂O₃ solution and dried over MgSO₄ and the solvents are then removed under reduced pressure. This gives 2.7 g of 4-iodomethyl-5-methyl-2-naphthalen-2-ylloxazole as a light-yellow solid. C₁₅H₁₂INO (349.17), MS(ESI): 350 (M+H⁺).

10

Methyl 2-(cis-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate



15

8.7 g of 1,3-cyclohexanediol and 12 g of dibutyltin oxide are dissolved in 600 ml of toluene and, in a water separator, heated under reflux. During the reaction, the reaction volume is reduced to half the original volume. After 4 hours, the reaction mixture is cooled to room temperature, and 300 ml of DMF, 9.0 g of methyl 2-bromomethyl-6-methylbenzoate and 9.4 g of cesium fluoride are added. The mixture is stirred at room temperature for 12 hours. The reaction mixture is diluted by addition of ethyl acetate and washed with saturated NaCl solution. The organic phase is dried over magnesium sulfate, the solvent is removed under reduced pressure and the residue is purified by flash chromatography on silica gel (n-heptane/ethyl acetate = 50:1 → 1:2). This gives 6 g of methyl 2-(cis-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate as an oil. C₁₆H₂₂O₄ (278.35), MS(ESI):

20


279 ($M + H^+$).

Methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate



8 g of methyl 2-(cis-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate are dissolved in 100 ml of vinyl acetate, and 1 g of *Candida antarctica* lipase B is added. The mixture is stirred at room temperature for seven hours and the enzyme is then filtered off and the solvent is removed under reduced pressure. The residue is purified by flash chromatography on silica gel (n-heptane/ethyl acetate = 10:1). This gives 3.9 g of the alcohol methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate as a colorless oil. $C_{16}H_{22}O_4$ (278.35), MS(ESI): 279 ($M + H^+$), ee = 98% (Chiralpak AD/2 250x4.6; n-heptane:ethanol:methanol = 25:1:0.5 + 0.1% trifluoroacetic acid, R_t = 8.9 min; retention time of the enantiomer: R_t = 9.9 min).

Methyl 2-methyl-6-[(1R,3S)-3-(5-methyl-2-naphthalen-2-yloxazol-4-ylmethoxy)-cyclohexyloxymethyl]benzoate




At room temperature, 50 mg of a 60% strength suspension of sodium hydride are added to a solution of 200 mg of methyl 2-((1R,3S)-3-hydroxycyclohexyloxy-

methyl)-6-methylbenzoate in 5 ml of dimethylformamide, and 380 mg of 4-iodomethyl-5-methyl-2-naphthalen-2-yloxazole are then added. After one hour, methyl tert-butyl ether is added and the mixture is extracted with water. The organic phase is dried over magnesium sulfate, the solvents are removed under reduced pressure and the residue is purified by RP-HPLC. This gives 94 mg of methyl 2-methyl-6-[(1R,3S)3-(5-methyl-2-naphthalen-2-yloxazol-4-ylmethoxy)cyclohexyloxymethyl]benzoate as a light-yellow oil.

C₃₁H₃₃NO₅ (499.61), MS(ESI): 500 (M + H⁺).

10

2-Methyl-6-[(1R,3S)3-(5-methyl-2-naphthalen-2-yloxazol-4-ylmethoxy)cyclohexyloxymethyl]benzoic acid



15


94 mg of methyl 2-methyl-6-[(1R,3S)3-(5-methyl-2-naphthalen-2-yloxazol-4-ylmethoxy)cyclohexyloxymethyl]benzoate are stirred at 90°C in a mixture of 10 ml of tert-butanol and 1 ml of 10 N potassium hydroxide solution. After two days, the mixture is acidified with hydrochloric acid and extracted with ethyl acetate. The combined organic phases are dried over magnesium sulfate, the solvents are removed under reduced pressure and the residue is purified by RP-HPLC. This gives 72 mg of 2-methyl-6-[(1R,3S)3-(5-methyl-2-naphthalen-2-yloxazol-4-ylmethoxy)cyclohexyloxymethyl]benzoic acid as an amorphous solid. C₃₀H₃₁NO₅ (485.59), MS(ESI): 486 (M + H⁺).

25

Example 2

Analogously to Example 1, diacetyl monoxime, benzo[1,3]dioxole-5-carbaldehyde


and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-[(1R,3S)-3-(2-benzof[1,3]dioxol-5-yl)-5-methyloxazol-4-ylmethoxy]cyclohexyloxymethyl]-6-methylbenzoic acid.



5 $C_{27}H_{29}NO_7$ (479.53), MS(ESI): 480 ($M + H^+$).

Example 3

Analogously to **Example 1**, diacetalmonoxime, 2,3-dihydrobenzo[1,4]dioxine-6-carbaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-[(1R,3S)-3-(2,3-dihydrobenzo[1,4]dioxin-6-yl)-5-methyloxazol-4-ylmethoxy]cyclohexyloxymethyl]-6-methylbenzoic acid.




15 $C_{28}H_{31}NO_7$ (493.56), MS(ESI): 494 ($M + H^+$).

Example 4

Analogously to **Example 1**, diacetyl monoxime, furan-2-carbaldehyde and methyl


2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-methyl-6-((1R,3S)-3-[5-methyl-2-(5-methylfuran-2-yl)oxazol-4-ylmethoxy]cyclohexyloxymethyl)benzoic acid.



5 $C_{25}H_{29}NO_6$ (439.51), MS(ESI): 440 ($M + H^+$).

Example 5

Analogously to Example 1, diacetyl monoxime, 5-methylthiophene-2-carbaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-methyl-6-((1R,3S)-3-[5-methyl-2-(5-methylthiophen-2-yl)oxazol-4-ylmethoxy]cyclohexyloxymethyl)benzoic acid.




15 $C_{25}H_{29}NO_5S$ (455.58), MS(ESI): 456 ($M + H^+$).

Example 6

Analogously to Example 1, diacetyl monoxime, 4-trifluoromethylsulfanylbenzaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-((1R,3S)-methyl-6-[3-[5-methyl-2-(4-trifluoromethylsulfanylphenyl)oxazol-4-ylmethoxy]cyclohexyloxymethyl]benzoic acid.


45



$C_{27}H_{28}F_3NO_5S$ (535.58), MS(ESI): 536 ($M + H^+$).

Example 7


- 5 Analogously to **Example 1**, diacetyl monoxime, 3-pentafluoroethoxybenzaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-((1R,3S)-methyl-6-(3-(5-methyl-2-[3-(1,1,2,2-tetrafluoroethoxy)phenyl]oxazol-4-ylmethoxy)cyclohexyloxymethyl)benzoic acid.



$C_{28}H_{29}F_4NO_6$ (551.54), MS(ESI): 552 ($M + H^+$).

Example 8


- 15 Analogously to **Example 1**, diacetyl monoxime, 4-phenoxybenzaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-((1R,3S)-methyl-6-(3-(5-methyl-2-(4-phenoxyphenyl)oxazol-4-ylmethoxy)cyclohexyloxymethyl)benzoic acid.



$C_{32}H_{33}NO_8$ (527.62), MS(ESI): 528 ($M + H^+$).

Example 9


- 5 Analogously to Example 1, diacetyl monoxime, thiophene-2-carbaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-((1R,3S)-methyl-6-[3-(5-methyl-2-thiophen-2-ylloxazol-4-ylmethoxy)cyclohexyl-oxymethyl]benzoic acid.





10 $C_{24}H_{27}NO_5S$ (441.55), MS(ESI): 442 ($M + H^+$).

Example 10

- 15 Analogously to Example 1, diacetyl monoxime, 3-fluoro-5-trifluoromethylbenzaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave methyl 2-((1R,3S)-[3-[2-(3-fluoro-5-trifluoromethylphenyl)-5-methyloxazol-4-ylmethoxy]cyclohexyloxymethyl]-6-methylbenzoate.



C₂₈H₂₉F₄NO₅ (535.54), MS(ESI): 536 (M + H⁺).




- 5 A mixture of 128 mg of methyl 2-((1R,3S)-3-[2-(3-fluoro-5-(trifluoromethyl)phenyl)-5-methyloxazol-4-ylmethoxy]cyclohexyloxymethyl)-6-methylbenzoate, 5 ml of ethylene glycol monomethyl ether and 0.6 ml of 10N KOH were heated under reflux for 24. After cooling, the mixture is acidified with hydrochloric acid and

extracted with ethyl acetate. The combined organic phases are dried over magnesium sulfate, the solvent is removed under reduced pressure and the residue is purified by RP-HPLC. This gives 56 mg of 2-((1R,3S)-3-(2-[3-(2-methoxyethoxy)-5-trifluoromethylphenyl]-5-methyloxazol-4-ylmethoxy)cyclohexyloxymethyl)-6-methylbenzoic acid as a colorless oil of molecular weight $C_{29}H_{32}F_3NO_7$ (563.58), MS(ESI): 564 ($M + H^+$).

Example 11

Analogously to Example 1, 1-phenyl-1,2-propanedione 2-oxime, p-tolualdehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-methyl-6-((1R,3S)-3-(5-phenyl-2-p-tolyloxazol-4-ylmethoxy)cyclohexyloxymethyl)benzoic acid.




$C_{32}H_{33}NO_5$ (511.62), MS(ESI) = 512 ($M + H^+$).

Example 12

Analogously to Example 1, 1-phenyl-1,2-propanedione 2-oxime, m-anisaldehyde and methyl 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-((1R,3S)-3-[2-(3-methoxyphenyl)-5-phenyloxazol-4-ylmethoxy]cyclohexyloxymethyl)-6-methylbenzoic acid.


49



$C_{32}H_{33}NO_6$ (527.62), MS(ESI) = 528 ($M+H^+$).

Example 13

Analogously to **Example 1**, 2-cyclohexyl-4-iodomethyloxazole and methyl
 5 2-((1R,3S)-3-hydroxycyclohexyloxymethyl)-6-methylbenzoate gave 2-((1R,3S)-3-
(2-cyclohexyloxazol-4-ylmethoxy)cyclohexyloxymethyl)-6-methylbenzoic acid.



$C_{25}H_{27}NO_5$ (421.50); MS(ESI): 422 ($M+H^+$).


10

We claim:

DEAV2003/0017

Dr. WI

1. A compound of the formula I



in which

10 Ring A is (C₃-C₈)-cycloalkanediyl or (C₃-C₈)-cycloalkenediyl, where in the cycloalkanediyl or cycloalkenediyl rings one or more carbon atoms may be replaced by oxygen atoms;

Ring B is a) phenyl; or

15 b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

20 R1 is a) in the case ring B = a):
SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

25 b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂, O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

c) in the case ring B = a) and R4 = phenyl:
(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

R2 is H or CF₃;

R4 is a) in the case ring B = a):
phenyl;

5

b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

10

c) in the case ring B = a) and R1 = a):
(C₁-C₆)-alkyl;

R5 is H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

R3 is H or (C₁-C₆)-alkyl;

15

X is (C₁-C₆)-alkanediyl, where in the alkanediyl group one or more carbon atoms may be replaced by oxygen atoms;

Y is (C₁-C₆)-alkanediyl, where in the alkanediyl group one or more carbon atoms may be replaced by oxygen atoms;

20

and its physiologically acceptable salts.

2. A compound of the formula I as claimed in claim 1, wherein

25

Ring A is (C₃-C₈)-cycloalkanediyl or (C₃-C₈)-cycloalkenediyl, where in the cycloalkanediyl or cycloalkenediyl rings one or more carbon atoms may be replaced by oxygen atoms;

30 Ring B is a) phenyl; or

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O

and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

R1 is

a) in the case ring B = a):

SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

5

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂,
O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-
alkyl;

10

c) in the case ring B = a) and R4 = phenyl:

(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

R is

H or CF₃;

15

R4 is

a) in the case ring B = a):
phenyl;

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

20

c) in the case ring B = a) and R1 = a):

(C₁-C₆)-alkyl;

25 R5 is

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

R3 is

H or (C₁-C₆)-alkyl;

X is

CH₂-O;

30

Y is

(C₁-C₆)-alkanediyl, where in the alkanediyl group one or more carbon
atoms may be replaced by oxygen atoms.

3. A compound of the formula I as claimed in claim 1 or 2, wherein

Ring A is (C₃-C₈)-cycloalkanediyl in which one carbon atom may be replaced by an oxygen atom;

Ring B is a) phenyl; or

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

R1 is a) in the case ring B = a):
SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂, O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

c) in the case ring B = a) and R4 = phenyl:
(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

R2 is H or CF₃;

R4 is a) in the case ring B = a):
phenyl;

b) in the case ring B = b):
H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

c) in the case ring B = a) and R1 = a):
(C₁-C₆)-alkyl;


R5 is H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

R3 is H or (C₁-C₈)-alkyl;

X is CH₂-O;

Y is CH₂-O.

4. A compound of the formula 1a



in which ring A, ring B, R1, R2, R3, R4, R5, X and Y are as defined in claims 1 to 3.

5. A compound of the formula 1a as claimed in claim 4 in which

R3 is H and

R5 is methyl.

6. A compound of the formula 1a as claimed in claim 4 or 5, in which

Ring A is (C₅-C₇)-cycloalkanedyl;

Ring B is a) phenyl; or

b) a 5- to 12-membered heteroaromatic ring which may contain one to four heteroatoms selected from the group consisting of N, O

and S, an 8- to 14-membered aromatic ring or (C₃-C₈)-cycloalkyl;

R1 is

a) in the case ring B = a):

SCF₃, OCF₂-CHF₂, O-phenyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-alkyl;

5

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, OCF₂-CF₃, SCF₃, OCF₂-CHF₂,
O-phenyl, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl-O-(C₁-C₃)-
alkyl;

10

c) in the case ring B = a) and R4 = phenyl:

(C₁-C₆)-alkyl or O-(C₁-C₆)-alkyl;

R2 is

H or CF₃;

15

R4 is

a) in the case ring B = a):

phenyl;

b) in the case ring B = b):

H, F, Cl, Br, OH, NO₂, CF₃, OCF₃, (C₁-C₆)-alkyl, O-(C₁-C₆)-alkyl;

20

c) in the case ring B = a) and R1/R2 = a):

(C₁-C₆)-alkyl;

25 R5 is

methyl;

R3 is

H;

X is

CH₂-O;

30

Y is

CH₂-O.

7. A compound of the formula I or Ia as claimed in any of claims 1 to 6,

wherein the central cycloalkanediyl ring is attached 1,3-cis.

8. A pharmaceutical comprising one or more compounds as claimed in one or more of claims 1 to 7.

5

9. A pharmaceutical comprising one or more compounds as claimed in one or more of claims 1 to 7 and one or more active substances which have favorable effects on metabolic disturbances or disorders frequently associated therewith.

10 10. A pharmaceutical comprising one or more compounds as claimed in one or more of claims 1 to 7 and one or more antidiabetics.

11. A pharmaceutical comprising one or more compounds as claimed in one or more of claims 1 to 7 and one or more lipid modulators.

15

12. The use of the compounds as claimed in one or more of claims 1 to 7 for the treatment and/or prevention of disorders of fatty acid metabolism and glucose utilization disorders.

20 13. The use of the compounds as claimed in one or more of claims 1 to 7 for the treatment and/or prevention of disorders in which insulin resistance is involved.

14. The use of the compounds as claimed in one or more of claims 1 to 7 for the treatment and/or prevention of diabetes mellitus including the prevention of the sequelae associated therewith.

25

15. The use of the compounds as claimed in one or more of claims 1 to 7 for the treatment and/or prevention of dyslipidemias and their sequelae.

30 16. The use of the compounds as claimed in one or more of claims 1 to 7 for the treatment and/or prevention of conditions which may be associated with the metabolic syndrome.

17. The use of compounds as claimed in one or more of claims 1 to 7 in combination with at least one further active compound for the treatment of disorders of fatty acid metabolism and glucose utilization disorders.
- 5 18. The use of compounds as claimed in one or more of claims 1 to 7 in combination with at least one further active compound for the treatment of disorders in which insulin resistance is involved.
- 10 19. A process for preparing a pharmaceutical comprising one or more of the compounds as claimed in one or more of claims 1 to 7, which comprises mixing the active compound with a pharmaceutically suitable carrier and bringing this mixture into a form suitable for administration.

Abstract

DEAV2003/0017

Dr.WI


Diarylcycloalkyl derivatives, process for their preparation and their use as pharmaceuticals

5

The invention relates to diarylcycloalkyl derivatives and to their physiologically acceptable salts, racemates and physiologically functional derivatives.

What is described are compounds of the formula I,

10



in which the radicals are as defined, and their physiologically acceptable salts and processes for their preparation. The compounds are suitable for the treatment and/or prevention of disorders of fatty acid metabolism and glucose utilization disorders as well as of disorders in which insulin resistance is involved.

15